



TITLE:

# The Remainder Term in the Dirichlet Divisor Problem (Analytic Number Theory)

AUTHOR(S):

TSANG, KAI-MAN

---

CITATION:

TSANG, KAI-MAN. The Remainder Term in the Dirichlet Divisor Problem (Analytic Number Theory). 数理解析研究所講究録 1996, 958: 111-119

ISSUE DATE:

1996-08

URL:

<http://hdl.handle.net/2433/60457>

RIGHT:

# The Remainder Term in the Dirichlet Divisor Problem

KAI-MAN TSANG

The University of Hong Kong  
Hong Kong

## 1. Introduction

Let  $d(n)$  denote the divisor function. We shall use  $c, c', c_1, c_2, \dots$  etc to denote certain constants which need not be the same at each occurrence. In this talk we shall give a survey on some recent results concerning the well-known remainder term

$$\Delta(x) := \sum_{n \leq x} d(n) - x(\log x + 2\gamma - 1), \quad x \geq 2,$$

which occurs frequently in analytic number theory. This can also be interpreted as a lattice point problem since  $\sum_{n \leq x} d(n)$  counts the number of lattice points in the first quadrant bounded by the hyperbola  $uv = x$ .

The first result on  $\Delta(x)$  was obtained more than one and a half century ago by Dirichlet, who proved by an elementary argument that  $\Delta(x) \ll \sqrt{x}$ . This upper bound was successively improved upon by many authors and the best result to date is :  $\Delta(x) \ll x^{\frac{7}{22} + \varepsilon}$  for any  $\varepsilon > 0$ , due to Iwaniec and Mozzochi [9]. It has been widely conjectured that  $\Delta(x) \ll_{\varepsilon} x^{1/4 + \varepsilon}$  is true for any  $\varepsilon > 0$ .

## 2. Values of $\Delta(x)$

Figure 1 below shows the graphs of  $y = \Delta(x)$  for four different ranges of  $x$ . An immediate observation was that  $\Delta(x)$  is highly oscillatory, it takes large values in both the positive and negative sides and yet it is slightly skewed towards the positive. Indeed, Voronoi [18] has proved in 1904 that

$$(1) \quad \int_2^X \Delta(x) dx = \frac{1}{4}X + \mathcal{O}(X^{3/4}),$$

that is,  $\Delta(x)$  has  $1/4$  as mean value. Concerning the large values of  $\Delta(x)$ , Hardy [3] showed earlier this century that

$$\Delta(x) = \begin{cases} \Omega_+ \left( (x \log x)^{\frac{1}{4}} \log \log x \right), \\ \Omega_- (x^{\frac{1}{4}}) \end{cases}$$

The best results in this direction to date are

$$\Delta(x) = \Omega_- \left\{ x^{\frac{1}{4}} \exp(c(\log \log x)^{\frac{1}{4}} (\log \log \log x)^{-\frac{3}{4}}) \right\}$$

and

$$\Delta(x) = \Omega_+ \left\{ (x \log x)^{\frac{1}{4}} (\log \log x)^{\frac{1}{4}(3+\log 4)} \exp(-c(\log \log \log x)^{\frac{1}{2}}) \right\}$$

for some constant  $c > 0$ , due to Corrádi - Kátai [1] and Selberg - Hafner [2] respectively. These  $\Omega$ -results, however, do not localize the occurrence of the extreme values of  $\Delta(x)$ . There is an earlier result of Tong [12] which says that :

*There exist positive constants  $c$  and  $c'$  such that, for any  $X \geq 1$  and for any  $t \in [-cX^{1/4}, cX^{1/4}]$  the equation  $\Delta(x) = t$  always has a solution  $x$  in the interval  $[X, X + c'\sqrt{X}]$ . In particular,  $\Delta(x)$  changes signs in  $[X, X + c'\sqrt{X}]$  for every  $X \geq 1$ , that is, the gap between the zeros of  $\Delta(x)$  is  $O(\sqrt{x})$ .*

Basing upon some numerical evidence, Ivić and te Riele [8] conjectured that  $\Delta(x)$  changes signs in every interval  $[X, X + c_\varepsilon X^{1/4+\varepsilon}]$  for any  $\varepsilon > 0$ ,  $X \geq X_0(\varepsilon)$  and  $c_\varepsilon$  is a constant dependent on  $\varepsilon$ . This conjecture, however, was shown to be too strong by the following result.

Heath-Brown and Tsang [5] : *There exist positive constants  $c, c_1, c_2$  such that, for any sufficiently large  $X$ , there are more than  $c_1 \sqrt{X} \log^5 X$  disjoint subintervals of length  $c_2 \sqrt{X} \log^{-5} X$  in  $[X, 2X]$ , throughout each of which either  $\Delta(x) > cX^{\frac{1}{4}}$  or  $\Delta(x) < -cX^{\frac{1}{4}}$  holds. In particular  $\Delta(x)$  does not change signs in each of these subintervals.*

The graph of  $y = \Delta(x)$  oscillates rigorously above and below the  $x$ -axis. Apparently there is no simple way to describe the values of  $\Delta(x)$ . However Heath-Brown [4] has shown that  $\Delta(x)$  possesses a distribution function in the following sense.

*There is a smooth function  $f(x)$  such that, for any interval  $I$ , we have*

$$X^{-1} \text{ meas } \{x \in [2, X] : x^{-1/4} \Delta(x) \in I\} \rightarrow \int_I f(\alpha) d\alpha$$

as  $X \rightarrow \infty$ .

### 3. Mean square of $\Delta(x)$

When considered in the mean, the remainder term  $\Delta(x)$  exhibits much better regularity. Voronoi's formula (1) shows that  $\Delta(x)$  has an asymptotic mean value of  $1/4$  over intervals of length  $\gg X^{3/4}$ . For the mean square, we have the following formula of Tong [13]:

$$\int_2^X \Delta(x)^2 dx = cX^{3/2} + F(X),$$

where  $c = (6\pi^2)^{-1} \sum_{n=1}^{\infty} d(n)^2 n^{-3/2} = 0.6542869 \dots$  and  $F(X) \ll X \log^5 X$ . Thus,  $\Delta(x)^2$  has asymptotic mean value of  $\frac{3}{2}c\sqrt{x}$  over intervals of length  $\gg \sqrt{X} \log^5 X$ . After more than thirty years have elapsed, this was then sharpened slightly by Preissmann [11] to  $F(X) \ll X \log^4 X$ , by using a variant of Hilbert's inequality. (Motohashi and others observed that the same can be obtained via an estimate for the sum  $\sum_{m \leq x} d(m)d(m+h)$ .)

There is not much information on the true order of  $F(X)$ . Ivić [6, Theorem 3.8] observed that

$$(2) \quad F(X) \ll U(X) \Rightarrow \Delta(x) \ll (U(x) \log x)^{1/3}.$$

Consequently, in view of the  $\Omega$ -results above, Ivić- Ouellet [7] showed that

$$F(x) = \Omega(x^{\frac{3}{4}}(\log x)^{-\frac{1}{4}}(\log \log x)^{\frac{3}{4}(3+\log 4)}e^{-c(\log \log \log x)^{1/2}}),$$

and it was even conjectured that  $F(x) \ll_{\varepsilon} x^{3/4+\varepsilon}$  is true for any  $\varepsilon > 0$ . This conjecture is very strong, since by Ivić's argument in (2), it implies the long standing conjecture that  $\Delta(x) \ll_{\varepsilon} x^{1/4+\varepsilon}$ . Ivić's conjecture is indeed too optimistic. Recently Tsang [15] deduced from the lower estimate :

$$(3) \quad \int_X^{2X} (F(x + \sqrt{X}) - F(x))^2 dx \gg X^3$$

that  $F(x) = \Omega(x)$ . Later, this was further sharpened by Lau - Tsang [17] to  $F(x) = \Omega_-(x \log^2 x)$ , which is an immediate consequence of the asymptotic formula

$$\int_2^X F(x) dx = -(8\pi^2)^{-1} X^2 \log^2 X + cX^2 \log X + \mathcal{O}(X^2).$$

This asymptotic formula can be reformulated as

$$\int_2^X (F(x) + (4\pi^2)^{-1} x \log^2 x - \kappa x \log x) dx \ll X^2$$

for a suitable constant  $\kappa$ . This leads us to the following

**Conjecture :**

$$(4) \quad F(x) = -(4\pi^2)^{-1}x \log^2 x + \kappa x \log x + \mathcal{O}(x),$$

that is,

$$(5) \quad \int_2^X \Delta(x)^2 dx = cX^{3/2} - (4\pi^2)^{-1}X \log^2 X + \kappa X \log X + \mathcal{O}(X).$$

This conjecture, if true, would imply that  $\Delta(x)^2$  has asymptotic mean value over intervals of length  $\gg \sqrt{X}$ . The  $\Omega$ -result for  $F(x)$  explains adequately the level of difficulty one faces in improving the upper bound for  $F(x)$ . The gap now left behind between the upper and lower bounds for  $F(x)$ , though small, seems very difficult to close.

Concerning the formula (4), one may naturally ask whether the  $\mathcal{O}(x)$  term contains some other main terms. Clearly

$$(6) \quad \int_X^{2X} (F(x + \sqrt{X}) - F(x)) dx = \int_{2X}^{2X + \sqrt{X}} F(x) dx - \int_X^{X + \sqrt{X}} F(x) dx \\ \ll X^{3/2} \log^4 X,$$

by applying Preissmann's upper bound for  $F(x)$ . We see that  $F(x + \sqrt{X}) - F(x)$  must change signs, for otherwise, by (6)

$$\int_X^{2X} (F(x + \sqrt{X}) - F(x))^2 dx \ll X \log^4 X \int_X^{2X} |F(x + \sqrt{X}) - F(x)| dx \ll X^{5/2} \log^8 X$$

which contradicts (3). Hence the term  $\mathcal{O}(x)$  in (4) is oscillatory and cannot be  $o(x)$ .

In support of our conjecture (4), we prove recently that :

Tsang [16]. *For some positive constant  $c$ , we have*

$$\int_2^X |F(x) + (4\pi^2)^{-1}x \log^2 x - \kappa x \log x|^r dx \ll (cr)^{4r} X^{r+1}$$

for any  $r \geq 1$ . Consequently, if  $H(x)$  is any increasing function satisfying  $2 \leq H(x) \leq \log^4 x$ , we have

$$|F(x) + (4\pi^2)^{-1}x \log x - \kappa x \log x| \leq xH(x)$$

for all but  $\mathcal{O}(Xe^{-cH(X)^{1/4}})$  values of  $x$  in  $[2, X]$ .

So at least our conjecture (4) is true for almost all  $x$ . Even though the constant  $\kappa$  is effectively computable, it is difficult to obtain an accurate numerical value for it. Very roughly, we estimated  $\kappa \doteq 0.32$  and Figure 2 below shows the graphs of  $y = \int_2^X \Delta(x)^2 dx - cX^{3/2} + (4\pi^2)^{-1}X \log^2 X - \kappa X \log X$  (with  $\kappa = 0.32$ ) for four different ranges of  $X$ .

In an earlier paper [14], we have established the following higher power moments of  $\Delta(x)$ :

$$\begin{aligned} \int_2^X \Delta(x)^3 dx &= c_3 X^{7/4} + \mathcal{O}(X^{7/4-\delta}), \\ \int_2^X \Delta(x)^4 dx &= c_4 X^2 + \mathcal{O}(X^{2-\delta}), \end{aligned}$$

where  $\delta$  is some small positive constant. Likewise we can consider the refinement of the  $\mathcal{O}$ -terms in these formulas, but the machinery available does not seem to be strong enough for this purpose.

#### 4. Conclusion

Our investigation on the error term  $\Delta(x)$  can be carried out for certain other error terms in number theory which have representation by Voronoi's type formula. These include  $E(T)$  defined by

$$E(T) := \int_0^T |\zeta(\tfrac{1}{2} + it)|^2 dt - T \log \frac{T}{2\pi} - (2\gamma - 1)T, \quad T \geq 2$$

and

$$P(x) = \sum_{n \leq x} r(n) - \pi x,$$

where  $r(n)$  denotes the number of integer pairs  $(x, y)$  such that  $n = x^2 + y^2$ . All our results on  $\Delta(x)$  hold true for  $E(T)$  and  $P(x)$ . The details will appear in a forthcoming paper.

After this talk was presented, Professor K. Matsumoto has kindly informed me that a conjectural formula of the shape

$$\int_0^T E(t)^2 dt \sim cT^{3/2} + c'T \log^A T$$


for a certain constant  $A$  has been proposed by him earlier [10]. Eventhough this is not as precise as our conjecture in (5), it is nonetheless in that same direction.

## REFERENCES

- [1] K. CORRÁDI-I. KÁTAI. *A comment on K.S. Gangadharan's paper entitled "Two classical lattice point problems"*. Magyar Tud. Akad. Mat. Fiz. Oszt. Kozl. 17, 89-97 (1967).
- [2] J.L. HAFNER. *New omega theorems for two classical lattice point problems*. Invent. Math. 63, 181-186 (1981).
- [3] G.H. HARDY. *On Dirichlet's divisor problem*. Proc. London Math. Soc. (2) 15, 1-25 (1916).
- [4] D.R. HEATH-BROWN. *The distribution and moments of the error term in the Dirichlet divisor problem*. Acta Arith. 60, 389-415 (1992).
- [5] D.R. HEATH-BROWN - K. TSANG. *Sign changes of  $E(T)$ ,  $\Delta(x)$  and  $P(x)$* . J. Number Theory, 49(1), 73-83 (1994).
- [6] A. IVIĆ. *Lectures on mean values of the Riemann zeta function*. Tata Inst. Fund. Research, Bombay, Springer-Verlag, Berlin, Heidelberg, New York, Tokyo (1991).
- [7] A. IVIĆ - M. OUELLET. *Some new estimates in the Dirichlet divisor problem*. Acta Arith. 52, 241-253 (1989).
- [8] A. IVIĆ - H.J.J. TE RIELE. *On the zeros of the error term for the mean square of  $|\zeta(\frac{1}{2} + it)|$* . Math. Comp. 56, 303-328 (1991).
- [9] H. IWANIEC - C.J. MOZZOCHI. *On the divisor and circle problems*. J. Number Theory, 29, 60-93 (1988).
- [10] K. MATSUMOTO. *Mean values of error terms in the theory of the Riemann zeta-function* (written in Japanese). Proc. Symposium Analytic Number Theory and Related Topics, Gakushuin Univ., 101-110 (1992).
- [11] E. PREISSMANN. *Sur la moyenne quadratique du terme de reste du problème du cercle*. C.R. Acad. Sci. Paris Sér. I 306, 151-154 (1988).
- [12] K.C. TONG. *On divisor problem I*. Acta Math. Sinica 5, (3), 313-324 (1955).
- [13] K.C. TONG. *On divisor problem III*. Acta Math. Sinica 6, (4), 515-541 (1956).
- [14] K. TSANG. *Higher-power moments of  $\Delta(x)$ ,  $E(t)$  and  $P(x)$* . Proc. London Math. Soc. (3) 65, 65-84 (1992).
- [15] K. TSANG. *A mean value theorem in the Dirichlet divisor problem*. preprint.
- [16] K. TSANG. *Mean square of the remainder term in the Dirichlet divisor problem II*. To appear in Acta Arith.
- [17] Y.K. LAU - K. TSANG. *Mean square of the remainder term in the Dirichlet*

- divisor problem I.* To appear in Journal de Théorie des Nombres de Bordeaux, Journées Arithmétiques (1993).
- [18] G.F. VORONOI. *Sur une fonction transcendante et ses applications à la sommation de quelques séries.* Ann. École Normale (3) 21 207-268, 459-534 (1904).








Figure 2. Graphs of  $y = \int_2^X \Delta(x)^2 dx - cX^{3/2} + (4\pi^2)^{-1} X \log^2 X - 0.32X \log X$